

Pneumomediastinum and Pneumothorax in a Neonate with Meconium Aspiration Syndrome

Pneumomediastino e Pneumotórax em Recém-Nascido com Síndrome de Aspiração Meconial

José Cunha de Alarcão^{1*}, Raquel S. Gonçalves¹, Mayara Nogueira², Ana Rodrigues Silva²

*Corresponding Author/Autor Correspondente:

José Miguel Cunha de Alarcão [josecunhaalarcao@gmail.com]
Rua O Conimbricense, nº66, 2ºA, 3030-504 Coimbra, Portugal
ORCID iD: 0000-0002-4649-9500

KEYWORDS: Infant, Newborn; Meconium Aspiration Syndrome/complications; Mediastinal Emphysema; Pneumothorax

PALAVRAS-CHAVE: Enfisema Mediastínico; Pneumotórax; Recém-Nascido; Síndrome de Aspiração Meconial/complicações

A term neonate born through meconium-stained amniotic fluid from instrumented delivery, (Apgar score 7/8), without need for resuscitation, presents with grunting, cyanosis and respiratory distress by the 5th minute of life, with pulse oximetry revealing a 70%-75% saturation which resolved with oxygenotherapy.

The neonate was admitted to the neonatal intensive care unit (NICU), requiring oxygenotherapy (maximum FiO₂ 40%, incubator environmental servo-oxygen delivery) for a saturation >95%. Initial bloodwork included a venous blood gases analysis (pH 7.26, pCO₂ 49.9 mmHg, HCO3- 22.4 mmol/L, BE -4.7 mmol/L, lactate 4.3 mmol/L), non-elevated C-reactive protein and a normal hemogram.

A chest x-ray showed a small pneumothorax (left lung apex) and left hemithorax radiolucency (outlining the mediastinal structures) with extrapleural air sign, sug-

gesting a large pneumomediastinum. On the right hemithorax, diffuse, asymmetric, patchy pulmonary opacities mixed with focal overinflation areas are suggestive of meconium aspiration syndrome (MAS) (Fig. 1).

Oxygen was successfully weaned off, with no signs of respiratory distress by the second day. A subsequent chest x-ray on the sixth day, showed complete reabsorption of the pneumomediastinum and pneumothorax (Fig. 2), after which the neonate was discharged from the NICU.

Pneumomediastinum is uncommon in neonates (2.5:1000 live births) but should be kept in mind particularly in cases of preterm neonates, meconium aspiration, positive pressure ventilation or congenital pneumonia.¹ Patients are usually asymptomatic, requiring no specific treatment, but should be monitored for evidence of cardio-respiratory compromise or development of other

1. Serviço de Pediatria, Hospital Pediátrico de Coimbra, CHUC, Coimbra, Portugal. 2. Serviço de Neonatologia, CHUC, Coimbra, Portugal.

Received/Recebido: 14/03/2022 - Accepted/Acete: 17/05/2022 - Published online/Publicado online: 02/06/2022 - Published/Publicado: 30/06/2022
© Author(s) (or their employer(s)) and Gazeta Médica 2022. Re-use permitted under CC BY-NC. No commercial re-use. © Autor (es) (ou seu (s) empregador (es)) e Gazeta Médica 2022. Reutilização permitida de acordo com CC BY-NC. Nenhuma reutilização comercial.



FIGURE 1. Chest x-ray at admission in NICU with a large pneumomediastinum (arrow) in the left hemithorax, with an extrapleural air sign (arrowhead) and a small linear pneumothorax (dotted arrow) on the left apex. Also visible are diffuse and asymmetric patchy opacities, mixed with focal areas of overinflation on the right pulmonary parenchyma, suggestive of MAS.




FIGURE 2. Chest x-ray at sixth day of life showing complete reabsorption of the pneumomediastinum and pneumothorax.

air leaks, especially pneumothorax. Infants with tension pneumomediastinum should be treated urgently with ultrasound-guided percutaneous drainage.

A chest x-ray should always be performed (lateral and anteroposterior views),² although recent studies suggest a lung ultrasound can also help confirm the diagnosis.³ In the presence of a high index of suspicion, a correct and early diagnostic can prevent iatrogenic damage such

as unnecessary insertion of a thoracic drain or prolonged antibiotic therapy.

AUTHORS CONTRIBUTION/ CONTRIBUIÇÃO AUTORAL

JCA and RSG: Wrote the manuscript

MN and ARS: Critically revised the manuscript

JCA e RSG: Escreveram o manuscrito

MN e ARS: Fizeram a revisão crítica do manuscrito

RESPONSABILIDADES ÉTICAS

CONFLITOS DE INTERESSE: Os autores declaram a inexistência de conflitos de interesse na realização do presente trabalho.

FONTES DE FINANCIAMENTO: Não existiram fontes externas de financiamento para a realização deste artigo.

CONFIDENCIALIDADE DOS DADOS: Os autores declaram ter seguido os protocolos da sua instituição acerca da publicação dos dados de doentes.

CONSENTIMENTO: Consentimento do doente para publicação obtido.

PROVENIÊNCIA E REVISÃO POR PARES: Não comissionado; revisão externa por pares.

ETHICAL DISCLOSURES

CONFLICTS OF INTEREST: The authors have no conflicts of interest to declare.

FINANCING SUPPORT: This work has not received any contribution, grant or scholarship.

CONFIDENTIALITY OF DATA: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

PATIENT CONSENT: Consent for publication was obtained.

PROVENANCE AND PEER REVIEW: Not commissioned; externally peer reviewed.

REFERENCES

1. Steele RW, Metz JR, Bass JW, DuBois JJ. Pneumothorax and pneumomediastinum in the newborn. Radiology. 1971;98:629-32. doi: 10.1148/98.3.629.
2. Raissaki M, Modatsou E, Hatzidaki E. Spontaneous pneumomediastinum in A term newborn: atypical radiographic and ct appearances. BJR | Case Rep. 2019;5:20180081.
3. Küng E, Habrina L, Berger A, Werther T, Aichhorn L. Diagnosing pneumomediastinum in a neonate using a lung ultrasound. Lancet. 2021;398:e13. doi: 10.1016/S0140-6736(21)01592-0.